arXiv:1606.02913v1 [math.CA] 9 Jun 2016

On the Fourier Transform of Bessel Functions over Complex Numbers - I: the Spherical Case

Zhi Qi

Abstract. In this note, we shall prove a formula for the Fourier transform of spherical Bessel functions over complex numbers, viewed as the complex analogue of the classical formulae of Hardy and Weber. The formula has strong representation theoretic motivations in the Waldspurger correspondence over the complex field.

1. Introduction

1.1. Representation Theoretic Motivations. It is observed in the article [**BM2**] of Baruch and Mao that several classical identities due to Weber and Hardy on Bessel functions can be used to realize the Shimura-Waldspurger correspondence between representations of $\mathrm{PGL}_2(\mathbb{R})$ and genuine representations of $\widetilde{\mathrm{SL}}_2(\mathbb{R})$, when the involved Bessel functions are attached to such representations in a certain way.

First, we have the following formula of Weber (see [**EMOT1**, 1.13 (25), 2.13 (27)] and [**Wat**, 13.3 (5)]),

$$\int_0^\infty \frac{1}{\sqrt{x}} J_\nu\left(4\pi\sqrt{x}\right) e\left(\pm xy\right) dx = \frac{1}{\sqrt{2y}} e\left(\mp\left(\frac{1}{2y} - \frac{1}{8}\nu - \frac{1}{8}\right)\right) J_{\frac{1}{2}\nu}\left(\frac{\pi}{y}\right), \tag{1.1}$$

for $y > 0$, where $e(x) = e^{2\pi i x}$ and $J_\nu(x)$ is the Bessel function of the first kind of order ν. This formula is valid when $\Re\mathrm{e}\,\nu > -1$. In other words, the Fourier transform of $x^{-\frac{1}{2}} J_\nu\left(4\pi x^{\frac{1}{2}}\right)$ is equal to $x^{-\frac{1}{2}} J_{\frac{1}{2}\nu}\left(\pi x^{-1}\right)$ up to an exponential factor. Second, the identity of Hardy is of the same nature, which involves the modified Bessel function K_ν of the second kind of order ν, (see [**EMOT1**, 1.13 (28), 2.13 (30)])

$$\begin{aligned}\int_0^\infty \frac{1}{\sqrt{x}} K_\nu\left(4\pi\sqrt{x}\right) e\left(\pm xy\right) dx = -\frac{\pi}{2\sin(\pi\nu)} \frac{1}{\sqrt{2y}} e\left(\pm\left(\frac{1}{2y} + \frac{1}{8}\right)\right) \\ \left(e\left(\pm\frac{1}{8}\nu\right) J_{\frac{1}{2}\nu}\left(\frac{\pi}{y}\right) - e\left(\mp\frac{1}{8}\nu\right) J_{-\frac{1}{2}\nu}\left(\frac{\pi}{y}\right)\right),\end{aligned} \tag{1.2}$$

with $|\Re\mathrm{e}\,\nu| < 1$.

Taking $\nu = 2k-1$ in (1.1), with k a positive integer, the Bessel function of order $2k-1$, respectively $k - \frac{1}{2}$, is attached to a discrete series representation of $\mathrm{PGL}_2(\mathbb{R})$, respectively

2010 *Mathematics Subject Classification.* 42B10, 33C10.

$\widetilde{\mathrm{SL}}_2(\mathbb{R})$. Thus, in this case, (1.1) should be interpreted as the local ingredient at the real place of the correspondence due to Shimura, Shintani and Waldspurger between cusp forms of weight $2k$ and cusp forms of weight $k+\frac{1}{2}$. Likewise, two formulae (1.1) and (1.2), with $\nu = 2it$ for t real or $\nu = 2t$ for $0 < t < \frac{1}{2}$, may be combined to realize the Shimura-Waldspurger correspondence for principal or complementary series representations. See [**BM2**].

In the framework of the relative trace formula of Jacquet, the theory in [**BM2**] along with the corresponding non-archimedean theory in [**BM1**] is used in [**BM3**] to produce a Waldspurger-type formula a *totally real* number field.

1.2. Statement of Results. In this note, we shall consider the Fourier transform of spherical Bessel functions over $\mathbb{C}$. Such Bessel functions are associated to spherical representations of $\mathrm{PGL}_2(\mathbb{C})$. According to [**Qi**, §2.7, 3.2], we define

$$\boldsymbol{J}_\mu(z) = \frac{2\pi^2}{\sin(2\pi\mu)}\left(J_{-2\mu}(4\pi\sqrt{z})\,J_{-2\mu}\big(4\pi\sqrt{\overline{z}}\big) - J_{2\mu}(4\pi\sqrt{z})\,J_{2\mu}\big(4\pi\sqrt{\overline{z}}\big)\right). \tag{1.3}$$

It is understood that the right hand side should be in its limit form when the order μ is a half integer. Moreover, the expression in (1.3) is independent of the choice of $\arg z$.

We shall prove the following analogue of the formulae of Weber and Hardy (1.1, 1.2).

Theorem 1.1. *Suppose that $|\Re\mathrm{e}\,\mu| < \frac{1}{2}$. We have the identity*

$$\int_0^{2\pi}\int_0^{\infty} \boldsymbol{J}_\mu\left(xe^{i\phi}\right) e(-2xy\cos(\phi+\theta))dxd\phi = \frac{1}{4y}e\left(\frac{\cos\theta}{y}\right)\boldsymbol{J}_{\frac{1}{2}\mu}\left(\frac{1}{16y^2e^{2i\theta}}\right), \tag{1.4}$$

with $y \in (0,\infty)$ and $\theta \in [0,2\pi)$.

The integral in (1.4) only converges as iterated double integral. It will therefore be more suitable to interpret the identity (1.4) in the sense of distributions. For this, we need some notation and definitions.

Define $\mathrm{Tr}(z) = z + \overline{z}$. Denote $\mathbb{C}^\times = \mathbb{C} \smallsetminus \{0\}$.

Let $\mathscr{S}(\mathbb{C})$ denote the space of Schwartz functions on $\mathbb{C}$, that is, smooth functions on $\mathbb{C}$ that rapidly decay at infinity along with all of their derivatives. If rapid decay also occurs at zero, then we say the functions are Schwartz functions on $\mathbb{C}^\times$, and the space of such functions is denoted by $\mathscr{S}(\mathbb{C}^\times)$.

The Fourier transform $\widehat{f}$ of a Schwartz function $f \in \mathscr{S}(\mathbb{C})$ is defined by

$$\widehat{f}(u) = \iint_{\mathbb{C}} f(z)e(-\mathrm{Tr}(uz))\,idz\wedge d\overline{z}.$$

We have $\widehat{\widehat{f}}(z) = f(-z)$.

Corollary 1.2. *We have*

$$\iint_{\mathbb{C}^\times} \boldsymbol{J}_\mu(z)\,\widehat{f}(z)\frac{idz\wedge d\overline{z}}{|z|} = \frac{1}{2}\iint_{\mathbb{C}^\times} e\left(\mathrm{Tr}\left(\frac{1}{2u}\right)\right)\boldsymbol{J}_{\frac{1}{2}\mu}\left(\frac{1}{16u^2}\right) f(u)\frac{idu\wedge d\overline{u}}{|u|}, \tag{1.5}$$

for $f \in \mathscr{S}(\mathbb{C})$, under the assumption $|\Re\mathrm{e}\,\mu| < \frac{1}{2}$. Furthermore, (1.5) *remains valid for all values of μ if one assumes $\widehat{f} \in \mathscr{S}(\mathbb{C}^\times)$.*

1.3. Remarks. The spherical Bessel functions for $\mathrm{SL}_2(\mathbb{C})$ were first discovered by Miatello and Wallach [**MW**] in the study of the spherical Kuznetsov trace formula for real semisimple groups of real rank one. The non-spherical Bessel functions for $\mathrm{SL}_2(\mathbb{C})$ were found by Bruggeman, Motohashi and Lokvenec-Guleska [**BM4, LG**] in generalizing the Kuznetsov formula to the non-spherical setting. In an entirely different way, they were recently rediscovered by the author [**Qi**] as the $\mathrm{GL}_2(\mathbb{C})$ example of the (fundamental) Bessel functions arising in the Voronoĭ summation formula for $\mathrm{GL}_n(\mathbb{C})$.

The identity (1.4) or (1.5) may be interpreted from the viewpoint of the Shimura-Waldspurger correspondence between spherical representations of $\mathrm{PGL}_2(\mathbb{C})$ and $\mathrm{SL}_2(\mathbb{C})$. In order to develop the spectral theory of the relative trace formula for the Waldspurger correspondence over the complex field and furthermore extend the Waldspurger formula to an arbitrary number field as in [**BM2, BM3**], we still need a formula similar to (1.4) or (1.5) for non-spherical Bessel functions for $\mathrm{PGL}_2(\mathbb{C})$. Such a formula however seems difficult to prove as the author can not find formulae for classical Bessel functions that would be helpful for treating non-spherical Bessel functions for $\mathrm{PGL}_2(\mathbb{C})$.

2. Preliminaries on Classical Bessel Functions

Let $J_\nu(z)$, $Y_\nu(z)$, $H_\nu^{(1,2)}(z)$ denote the three kinds of Bessel functions of order ν and $I_\nu(z)$ the modified Bessel function of the first kind of oder ν.

$J_\nu(z)$ is defined by the series (see [**Wat**, 3.1 (8)])

$$J_\nu(z) = \sum_{n=0}^{\infty} \frac{(-)^n \left(\frac{1}{2}z\right)^{\nu+2n}}{n!\Gamma(\nu+n+1)}. \tag{2.1}$$

We have the following connection formulae (see [**Wat**, 3.61 (1, 2, 3, 4), 3.1 (8), 3.7 (2)])

$$J_\nu(z) = \frac{H_\nu^{(1)}(z) + H_\nu^{(2)}(z)}{2}, \qquad J_{-\nu}(z) = \frac{e^{\pi i\nu}H_\nu^{(1)}(z) + e^{-\pi i\nu}H_\nu^{(2)}(z)}{2}, \tag{2.2}$$

$$Y_\nu(z) = \frac{J_\nu(z)\cos(\pi\nu) - J_{-\nu}(z)}{\sin(\pi\nu)}, \quad Y_{-\nu}(z) = \frac{J_\nu(z) - J_{-\nu}(z)\cos(\pi\nu)}{\sin(\pi\nu)}, \tag{2.3}$$

$$I_\nu\left(e^{\pm\frac{1}{2}\pi i}z\right) = e^{\pm\frac{1}{2}\pi i\nu}J_\nu(z). \tag{2.4}$$

Let N be a nonnegative integer. We have the asymptotic expansions (see [**Wat**, 7.2 (1, 2)]),

$$H_\nu^{(1)}(z) = \left(\frac{2}{\pi z}\right)^{\frac{1}{2}} e^{i\left(z-\frac{1}{2}\nu\pi-\frac{1}{4}\pi\right)}\left(\sum_{n=0}^{N-1}\frac{\left(\frac{1}{2}-\nu\right)_n\left(\frac{1}{2}+\nu\right)_n}{n!(2iz)^n} + O\left(|z|^{-N}\right)\right), \tag{2.5}$$

$$H_\nu^{(2)}(z) = \left(\frac{2}{\pi z}\right)^{\frac{1}{2}} e^{-i\left(z-\frac{1}{2}\nu\pi-\frac{1}{4}\pi\right)}\left(\sum_{n=0}^{N-1}\frac{\left(\frac{1}{2}-\nu\right)_n\left(\frac{1}{2}+\nu\right)_n}{n!(-2iz)^n} + O\left(|z|^{-N}\right)\right), \tag{2.6}$$

of which (2.5) is valid when z is such that $-\pi+2\delta \leqslant \arg z \leqslant 2\pi-2\delta$, and (2.6) when $-2\pi+2\delta \leqslant \arg z \leqslant \pi-2\delta$, δ being any positive acute angle. Here, $(\lambda)_n = \prod_{k=0}^{n-1}(\lambda+k)$.

3. Two Formulae for Classical Bessel Functions

In this section, we shall give two formulae for classical Bessel functions that will be used for the proof of Theorem 1.1.

3.1. We have Weber's second exponential integral formula [**Wat**, 13.31 (1)]

$$\int_0^\infty J_\nu(ax)J_\nu(bx)\exp\left(-p^2x^2\right)xdx=\frac{1}{2p^2}\exp\left(-\frac{a^2+b^2}{4p^2}\right)I_\nu\left(\frac{ab}{2p^2}\right). \tag{3.1}$$

It is required that $\Re\mathrm{e}\,\nu>-1$ and $|\arg p|<\frac{1}{4}\pi$ to secure absolute convergence, but a and b are unrestricted nonzero complex numbers. It follows that

$$\begin{aligned}\int_0^\infty \left(J_{-\nu}\left(a\sqrt{x}\right)J_{-\nu}\left(\overline{a}\sqrt{x}\right)-J_\nu\left(a\sqrt{x}\right)J_\nu\left(\overline{a}\sqrt{x}\right)\right)\exp\left(-p^2x\right)dx\\ =\frac{1}{p^2}\exp\left(-\frac{a^2+\overline{a}^2}{4p^2}\right)\left(I_{-\nu}\left(\frac{|a|^2}{2p^2}\right)-I_\nu\left(\frac{|a|^2}{2p^2}\right)\right),\end{aligned} \tag{3.2}$$

if $|\Re\mathrm{e}\,\nu|<1$, $|\arg p|<\frac{1}{4}\pi$ and $a\neq 0$. We claim that (3.2) is still valid even if $\arg p=\pm\frac{1}{4}\pi$ in the sense that the left hand side of (3.2) remains convergent and converges to the value on the right.

Since the expressions on both sides of (3.2) are indeed independent on the argument of a modulo π, we shall assume that $-\frac{1}{2}\pi<\arg a\leqslant\frac{1}{2}\pi$.

To prove the convergence, we first partition the integral in (3.2) into two integrals over the intervals $(0,1]$ and $[1,\infty)$ respectively. Since $|\Re\mathrm{e}\,\nu|<1$, the first integral is absolutely convergent due to the series expansions of $J_\nu(z)$ and $J_{-\nu}(z)$ at zero (see (2.1)). As for the convergence of the second integral, using the connection formulae (2.2), we write

$$\begin{aligned}&J_{-\nu}\left(a\sqrt{x}\right)J_{-\nu}\left(\overline{a}\sqrt{x}\right)-J_\nu\left(a\sqrt{x}\right)J_\nu\left(\overline{a}\sqrt{x}\right)\\ &\quad=\frac{i\sin(\pi\nu)}{2}\left(e^{\pi i\nu}H_\nu^{(1)}\left(a\sqrt{x}\right)H_\nu^{(1)}\left(\overline{a}\sqrt{x}\right)-e^{-\pi i\nu}H_\nu^{(2)}\left(a\sqrt{x}\right)H_\nu^{(2)}\left(\overline{a}\sqrt{x}\right)\right).\end{aligned}$$

Therefore, in view of the following asymptotic of $H_\nu^{(1,2)}(z)$ at infinity (see (2.5) and (2.6)),

$$H_\nu^{(1,2)}(z)=\left(\frac{2}{\pi z}\right)^{\frac{1}{2}}\exp\left(\pm i\left(z-\frac{\pi\nu}{2}-\frac{\pi}{4}\right)\right)\left(1\pm\frac{1-4\nu^2}{8iz}+O\left(|z|^{-2}\right)\right),$$

with $|\arg z|\leqslant\frac{1}{2}\pi$, we are then reduced to the convergence of the integrals

$$\int_1^\infty\frac{1}{\sqrt{x}}\exp\left(\pm i(a+\overline{a})\sqrt{x}-p^2x\right)dx,\quad \int_1^\infty\frac{1}{x}\exp\left(\pm i(a+\overline{a})\sqrt{x}-p^2x\right)dx.$$

By partial integration, the former turns into

$$\begin{aligned}&\left(\frac{1}{p^2}\pm\frac{i(a+\overline{a})}{2p^4}\right)\exp\left(\pm i(a+\overline{a})-p^2\right)\\ &-\left(\frac{1}{2p^2}+\frac{(a+\overline{a})^2}{4p^4}\right)\int_1^\infty x^{-\frac{3}{2}}\exp\left(\pm i(a+\overline{a})\sqrt{x}-p^2x\right)dx\\ &\mp\frac{i(a+\overline{a})}{2p^4}\int_1^\infty x^{-2}\exp\left(\pm i(a+\overline{a})\sqrt{x}-p^2x\right)dx,\end{aligned}$$

in which the integrals converge absolutely whenever $|\arg p| \leqslant \frac{1}{4}\pi$. The convergence of the latter may be proven in the same way. This completes the proof of the convergence.

With the above arguments, it is easy to verify that the left hand side of (3.2) gives rise to a continuous function on the sector $\left\{p : |\arg p| \leqslant \frac{1}{4}\pi\right\}$. Then follows the second assertion in the claim.

Rescaling a by the factor 4π, letting $p^2 = 2\pi e^{\mp\frac{1}{2}\pi i}c$ and using the identity (2.4) to turn the $I_{\pm\nu}$ into $J_{\pm\nu}$, the claim above yields the following lemma.

LEMMA 3.1. *Suppose that* $|\Re\mathrm{e}\,\nu| < 1$, $a \neq 0$ *and* $c > 0$. *Then*

$$\int_0^\infty \left(J_{-\nu}\left(4\pi a\sqrt{x}\right)J_{-\nu}\left(4\pi\overline{a}\sqrt{x}\right) - J_\nu\left(4\pi a\sqrt{x}\right)J_\nu\left(4\pi\overline{a}\sqrt{x}\right)\right)e\left(\pm cx\right)dx$$
$$= \mp\frac{1}{2\pi i c}e\left(\mp\frac{a^2+\overline{a}^2}{c}\right)\left(e^{\mp\frac{1}{2}\pi i\nu}J_{-\nu}\left(\frac{4\pi|a|^2}{c}\right) - e^{\pm\frac{1}{2}\pi i\nu}J_\nu\left(\frac{4\pi|a|^2}{c}\right)\right).$$

3.2. According to [**EMOT2**, 8.7 (22)], we have the following formula

$$\int_0^1 J_\nu(ax)\frac{\sin\left(b\sqrt{1-x^2}\right)}{\sqrt{1-x^2}}dx + \int_1^\infty J_\nu(ax)\frac{\exp\left(-b\sqrt{x^2-1}\right)}{\sqrt{x^2-1}}dx$$
$$= \frac{\pi}{2}J_{\frac{1}{2}\nu}\left(\frac{\sqrt{a^2+b^2}-b}{2}\right)Y_{\frac{1}{2}\nu}\left(\frac{\sqrt{a^2+b^2}+b}{2}\right). \tag{3.3}$$

It is assumed in [**EMOT2**] that $\Re\mathrm{e}\,\nu > -1$ and $a, b > 0$. We shall prove however that (3.3) is valid even if $\Re\mathrm{e}\,b \geqslant 0$, provided that $\Re\mathrm{e}\,\nu > -1$ and $a > 0$.

Given $\Re\mathrm{e}\,\nu > -1$ and $a > 0$. First of all, the first integral on the left hand side of (3.3) is absolutely convergent for any complex number b, and, if one makes the additional assumption $\Re\mathrm{e}\,b \geqslant 0$, then so is the second. The formula (3.3) is a relation connecting functions of b which are analytic on the right half plane $\{b : \Re\mathrm{e}\,b > 0\}$ and extend continuously onto the imaginary axis $\{b : \Re\mathrm{e}\,b = 0\}$, where the principal branch of the square root is assumed on the right hand side. Therefore, by the principle of analytic continuation, (3.3) remains valid whenever $\Re\mathrm{e}\,b \geqslant 0$.

We shall be particularly interested in the case when $\Re\mathrm{e}\,b = 0$ and $|\Im\mathrm{m}\,b| \leqslant a$. With the observation that the first integral in (3.3) is odd with respect to b, along with the identities in (2.3) that connect $Y_{\pm\frac{1}{2}\nu}$ with $J_{\frac{1}{2}\nu}$ and $J_{-\frac{1}{2}\nu}$, we obtain the following lemma.

LEMMA 3.2. *Suppose that* $|\Re\mathrm{e}\,\nu| < 1$, $a > 0$ *and* $-a \leqslant c \leqslant a$. *If we put* $w = \frac{1}{2}\left(\sqrt{a^2-c^2}+ic\right)$, *then*

$$\int_1^\infty \left(J_{-\nu}(ax) + J_\nu(ax)\right)\frac{\cos\left(c\sqrt{x^2-1}\right)}{\sqrt{x^2-1}}dx$$
$$= \frac{\pi\cot\left(\frac{1}{2}\pi\nu\right)}{2}\left(J_{\frac{1}{2}\nu}(w)\,J_{\frac{1}{2}\nu}(\overline{w}) - J_{-\frac{1}{2}\nu}(w)\,J_{-\frac{1}{2}\nu}(\overline{w})\right).$$

4. Proof of Theorem 1.1

With the preparations in the last section, we are now ready to prove Theorem 1.1. Indeed, we shall prove the following reformulation of Theorem 1.1.

PROPOSITION 4.1. *Suppose that* $|\Re\mathfrak{e}\,\mu| < \frac{1}{2}$. *The iterated double integral*

$$\int_0^{2\pi}\int_0^{\infty}\Big(J_{-2\mu}\big(4\pi e^{\frac{1}{2}i\phi}\sqrt{x}\big)J_{-2\mu}\big(4\pi e^{-\frac{1}{2}i\phi}\sqrt{x}\big)- J_{2\mu}\big(4\pi e^{\frac{1}{2}i\phi}\sqrt{x}\big)J_{2\mu}\big(4\pi e^{-\frac{1}{2}i\phi}\sqrt{x}\big)\Big)e(\pm 2xy\cos(\phi+\theta))dx\,d\phi, \tag{4.1}$$

with $y\in(0,\infty)$ *and* $\theta\in\big(-\frac{1}{2}\pi,\frac{1}{2}\pi\big]$, *is equal to*

$$\frac{\cos(\pi\mu)}{2y}e\left(\mp\frac{\cos\theta}{y}\right)\left(J_{-\mu}\left(\frac{\pi}{ye^{i\theta}}\right)J_{-\mu}\left(\frac{\pi}{ye^{-i\theta}}\right)-J_{\mu}\left(\frac{\pi}{ye^{i\theta}}\right)J_{\mu}\left(\frac{\pi}{ye^{-i\theta}}\right)\right).$$

PROOF. If we substitute ϕ by $\phi-\theta$, partition the domain of ϕ into four open intervals $\big(0,\frac{1}{2}\pi\big)$, $\big(\frac{1}{2}\pi,\pi\big)$, $\big(\pi,\frac{3}{2}\pi\big)$ and $\big(\frac{3}{2}\pi,2\pi\big)$, and make a suitable change of variables for each resulting integral, then the integral (4.1) turns into the sum of four similar integrals, the first of which is

$$\int_0^{\frac{1}{2}\pi}\int_0^{\infty}\Big(J_{-2\mu}\big(4\pi e^{\frac{1}{2}i(\phi-\theta)}\sqrt{x}\big)J_{-2\mu}\big(4\pi e^{-\frac{1}{2}i(\phi-\theta)}\sqrt{x}\big)- J_{2\mu}\big(4\pi e^{\frac{1}{2}i(\phi-\theta)}\sqrt{x}\big)J_{2\mu}\big(4\pi e^{-\frac{1}{2}i(\phi-\theta)}\sqrt{x}\big)\Big)e(\pm 2xy\cos\phi)dx\,d\phi. \tag{4.2}$$

For $\phi\in\big(0,\frac{1}{2}\pi\big)$, upon choosing $\nu=2\mu$, $a=e^{\frac{1}{2}i(\phi-\theta)}$, $c=2y\cos\phi$ in Lemma 3.1, we deduce that the integral (4.2) is equal to

$$\begin{aligned}&\mp\frac{1}{4\pi iy}\int_0^{\frac{1}{2}\pi}\frac{1}{\cos\phi}e\left(\mp\frac{\cos(\phi-\theta)}{y\cos\phi}\right)\left(e^{\pm\pi i\mu}J_{-2\mu}\left(\frac{2\pi}{y\cos\phi}\right)-e^{\mp\pi i\mu}J_{2\mu}\left(\frac{2\pi}{y\cos\phi}\right)\right)d\phi\\ &=\mp\frac{1}{4\pi iy}e\left(\mp\frac{\cos\theta}{y}\right)\int_0^{\frac{1}{2}\pi}\frac{1}{\cos\phi}e\left(\mp\frac{\sin\theta\tan\phi}{y}\right)\\ &\qquad\left(e^{\pm\pi i\mu}J_{-2\mu}\left(\frac{2\pi}{y\cos\phi}\right)-e^{\mp\pi i\mu}J_{2\mu}\left(\frac{2\pi}{y\cos\phi}\right)\right)d\phi.\end{aligned}$$

The other three integrals that are similar to (4.2) can be computed in the same way. Summing these up, the integral (4.1) turns into

$$-\frac{\sin(\pi\mu)}{\pi y}e\left(\mp\frac{\cos\theta}{y}\right)\int_0^{\frac{1}{2}\pi}\frac{1}{\cos\phi}\cos\left(\frac{2\pi\sin\theta\tan\phi}{y}\right)\left(J_{-2\mu}\left(\frac{2\pi}{y\cos\phi}\right)+J_{2\mu}\left(\frac{2\pi}{y\cos\phi}\right)\right)d\phi. \tag{4.3}$$

Making the change of variables $t=1/\cos\phi$, the integral in (4.3) turns into

$$\int_1^{\infty}\left(J_{-2\mu}\left(\frac{2\pi}{y}t\right)+J_{2\mu}\left(\frac{2\pi}{y}t\right)\right)\cos\left(\frac{2\pi\sin\theta}{y}\sqrt{t^2-1}\right)\frac{dt}{\sqrt{t^2-1}}.$$

Applying Lemma 3.2 with $\nu=2\mu$, $a=2\pi/y$, $c=2\pi\sin\theta/y$ and $w=\pi e^{i\theta}/y$, the integral above is then equal to

$$\frac{\pi\cot(\pi\mu)}{2}\left(J_{\mu}\left(\frac{\pi}{ye^{i\theta}}\right)J_{\mu}\left(\frac{\pi}{ye^{-i\theta}}\right)-J_{-\mu}\left(\frac{\pi}{ye^{i\theta}}\right)J_{-\mu}\left(\frac{\pi}{ye^{-i\theta}}\right)\right).$$

This completes the proof. Q.E.D.

5. Proof of Corollary 1.2

We first make some remarks on the convergence of (1.5) for $f \in \mathscr{S}(\mathbb{C})$. In view of the asymptotics of J_v at zero and infinity, the right hand side of (1.5) absolutely converges for all μ, but the absolute convergence of left hand side only holds for $|\Re\mathfrak{e}\,\mu| < \frac{1}{2}$. We also have compact convergence and hence analyticity with respect to the order μ.

To prove (1.5) for $f \in \mathscr{S}(\mathbb{C})$ in Corollary 1.2, it suffices to verify the identity,

$$\begin{aligned}\int_0^{2\pi}\int_0^{\infty}\int_0^{2\pi}\int_0^{\infty} \boldsymbol{J}_\mu\left(xe^{i\phi}\right) e(-2xy\cos(\phi+\theta))\, f\left(ye^{i\theta}\right) y\,dy\,d\theta\,dx\,d\phi \\ = \frac{1}{4}\int_0^{2\pi}\int_0^{\infty} e\left(\frac{\cos\theta}{y}\right) \boldsymbol{J}_{\frac{1}{2}\mu}\left(\frac{1}{16y^2e^{2i\theta}}\right) f\left(ye^{i\theta}\right) dy\,d\theta.\end{aligned} \tag{5.1}$$

Since the idea is straightforward, we shall only give a sketch of the proof as below and leave the details to the readers. Starting from the integral on the left hand side of (5.1), we write the inner integral in the Cartesian coordinates and apply the combination of partial integrations that have the effect of dividing $-16\pi^2x^2-1$. After this, we change the order of integrations, which is legitimate since $\boldsymbol{J}_\mu\left(xe^{i\phi}\right)/(16\pi^2x^2+1)$ is absolutely integrable. Then, we apply the partial integrations reverse to those that we performed at the beginning and rewrite the outer integral in the polar coordinates. In this ways, we retrieve the expression on the left hand side of (5.1) but with changed integration order. Here, to prove that some differentiations under the integral sign are permissible, we need to also verify the compact convergence of certain integrals, especially the integral in Lemma 3.1, with respect to parameters. Finally, integrating out the resulted inner integral using (1.4), we arrive at the integral on the right hand side of (5.1).

When $\widehat{f} \in \mathscr{S}(\mathbb{C}^\times)$, the integral on the left hand side of (1.5) becomes absolutely convergent for all μ, then follows the second assertion in Corollary 1.2.

References

[BM1] Ehud Moshe Baruch and Zhengyu Mao. Bessel identities in the Waldspurger correspondence over a p-adic field. *Amer. J. Math.*, 125(2):225–288, 2003.

[BM2] Ehud Moshe Baruch and Zhengyu Mao. Bessel identities in the Waldspurger correspondence over the real numbers. *Israel J. Math.*, 145:1–81, 2005.

[BM3] Ehud Moshe Baruch and Zhengyu Mao. Central value of automorphic L-functions. *Geom. Funct. Anal.*, 17(2):333–384, 2007.

[BM4] Roelof W. Bruggeman and Yoichi Motohashi. Sum formula for Kloosterman sums and fourth moment of the Dedekind zeta-function over the Gaussian number field. *Funct. Approx. Comment. Math.*, 31:23–92, 2003.

[EMOT1] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Tables of integral transforms. Vol. I.* McGraw-Hill Book Company, Inc., New York-Toronto-London, 1954. Based, in part, on notes left by Harry Bateman.

[EMOT2] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Tables of integral transforms. Vol. II.* McGraw-Hill Book Company, Inc., New York-Toronto-London, 1954. Based, in part, on notes left by Harry Bateman.

[LG] Hristina Lokvenec-Guleska. *Sum Formula for SL_2 over Imaginary Quadratic Number Fields*. Ph.D. Thesis. Utrecht University, 2004.

[MW] R. Miatello and N. R. Wallach. Kuznetsov formulas for real rank one groups. *J. Funct. Anal.*, 93(1):171–206, 1990.

[Qi] Zhi Qi. *Theory of Bessel functions of high rank*. Ph.D. Thesis. The Ohio State University, 2015.

[Wat] G. N. Watson. *A Treatise on the Theory of Bessel Functions*. Cambridge University Press, Cambridge, England; The Macmillan Company, New York, 1944.

Department of Mathematics, Rutgers University, Hill Center, 110 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA

E-mail address: zhi.qi@rutgers.edu